

## CLAIMS

### WHAT IS CLAIMED IS:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group  
5 consisting of SEQ ID NO: 1 – 10, a mature protein coding portion of SEQ ID NO: 1 – 10, an  
active domain of SEQ ID NO: 1 – 10, and complementary sequences thereof.
2. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said  
10 polynucleotide hybridizes to the polynucleotide of claim 1 under stringent hybridization  
conditions.
3. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said  
polynucleotide has greater than about 90% sequence identity with the polynucleotide of claim 1.
4. The polynucleotide of claim 1 wherein said polynucleotide is DNA.
5. An isolated polynucleotide of claim 1 wherein said polynucleotide comprises the  
complementary sequences.
6. A vector comprising the polynucleotide of claim 1.
7. An expression vector comprising the polynucleotide of claim 1.
8. A host cell genetically engineered to comprise the polynucleotide of claim 1.  
25
9. A host cell genetically engineered to comprise the polynucleotide of claim 1 operatively  
associated with a regulatory sequence that modulates expression of the polynucleotide in the host  
cell.
- 30 10. An isolated polypeptide, wherein the polypeptide is selected from the group consisting  
of:
  - a) a polypeptide encoded by any one of the polynucleotides of claim 1; and

b) a polypeptide encoded by a polynucleotide hybridizing under stringent conditions with any one of SEQ ID NO: 1 – 10.

11. A composition comprising the polypeptide of claim 10 and a carrier.

12. An antibody directed against the polypeptide of claim 10.

13. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

a) contacting the sample with a compound that binds to and forms a complex with the polynucleotide of claim 1 for a period sufficient to form the complex; and

b) detecting the complex, so that if a complex is detected, the polynucleotide of claim 1 is detected.

14. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

a) contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 1 under such conditions;

b) amplifying a product comprising at least a portion of the polynucleotide of claim 1; and

c) detecting said product and thereby the polynucleotide of claim 1 in the sample.

15. The method of claim 14, wherein the polynucleotide is an RNA molecule and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.

16. A method for detecting the polypeptide of claim 10 in a sample, comprising:

a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and

b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 10 is detected.

17. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

- a) contacting the compound with the polypeptide of claim 10 under conditions sufficient to form a polypeptide/compound complex; and
- b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

5

18. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

- a) contacting the compound with the polypeptide of claim 10, in a cell, under conditions sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and
- b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500

19. A method of producing the polypeptide of claim 10, comprising,

- a) culturing a host cell comprising a polynucleotide sequence selected from the group consisting of a polynucleotide sequence of SEQ ID NO: 1-10, a mature protein coding portion of SEQ ID NO: 1-10, an active domain of SEQ ID NO: 1-10, complementary sequences thereof and a polynucleotide sequence hybridizing under stringent conditions to SEQ ID NO: 1-10, under conditions sufficient to express the polypeptide in said cell; and
- b) isolating the polypeptide from the cell culture or cells of step (a).

20. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of any one of the polypeptides from the Sequence Listing, the mature protein portion thereof, or the active domain thereof.

21. The polypeptide of claim 20 wherein the polypeptide is provided on a polypeptide array.

22. A collection of polynucleotides, wherein the collection comprising the sequence information of at least one of SEQ ID NO: 1 – 10.

30

23. The collection of claim 22, wherein the collection is provided on a nucleic acid array.

24. The collection of claim 23, wherein the array detects full-matches to any one of the polynucleotides in the collection.

5

25. The collection of claim 23, wherein the array detects mismatches to any one of the polynucleotides in the collection.

10

26. The collection of claim 22, wherein the collection is provided in a computer-readable format.

27. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

15

28. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising an antibody that specifically binds to a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

20

25

30

35

40

45

50

55

60

65

70

75

80

85

90

95